



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Genome sequence of the filamentous bacterium *Fibrisoma limi* BUZ 3T

Filippini, Manuela ; Qi, Weihong ; Jaenicke, Sebastian ; Goesmann, Alexander ; Smits, Theo H M ;
Bagheri, Homayoun C

Abstract: *Fibrisoma limi* strain BUZ 3T, a Gram-negative bacterium, was isolated from coastal mud from the North Sea (Fedderwardersiel, Germany) and characterized using a polyphasic approach in 2011. The genome consists of a chromosome of about 7.5 Mb and three plasmids.

DOI: <https://doi.org/10.1128/JB.00869-12>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-77617>

Journal Article

Published Version

Originally published at:

Filippini, Manuela; Qi, Weihong; Jaenicke, Sebastian; Goesmann, Alexander; Smits, Theo H M; Bagheri, Homayoun C (2012). Genome sequence of the filamentous bacterium *Fibrisoma limi* BUZ 3T. *Journal of Bacteriology*, 16(194):4445.

DOI: <https://doi.org/10.1128/JB.00869-12>

Updated information and services can be found at:
<http://jb.asm.org/content/194/16/4445>

REFERENCES

These include:

This article cites 4 articles, 3 of which can be accessed free at:
<http://jb.asm.org/content/194/16/4445#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Genome Sequence of the Filamentous Bacterium *Fibrisoma limi* BUZ 3^T

Manuela Filippini,^a Weihong Qi,^b Sebastian Jaenicke,^c Alexander Goesmann,^c Theo H. M. Smits,^{d,e} and Homayoun C. Bagheri^a

Institute of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland^a; Functional Genomics Center Zurich, Uni/ETH Zurich, Zurich, Switzerland^b; CeBiTec, University of Bielefeld, Bielefeld, Germany^c; Agroscope Changins-Wädenswil ACW, Division of Plant Protection, Wädenswil, Switzerland^d; and Smits Bioinformatics Consultancy, Samstagern, Switzerland^e

***Fibrisoma limi* strain BUZ 3^T, a Gram-negative bacterium, was isolated from coastal mud from the North Sea (Fedderwardersiel, Germany) and characterized using a polyphasic approach in 2011. The genome consists of a chromosome of about 7.5 Mb and three plasmids.**

Fibrisoma limi BUZ 3^T is a member of the genus *Fibrisoma* (*Bacteroidetes*, *Sphingobacteriales*, *Cytophagaceae*), was characterized in 2011 (1), and produces multicellular filaments during growth, reaching up to 90 μm in length after 1 day of growth in liquid culture. This bacterium does not produce flagella, and gliding motility was not observed.

Bacteria were grown overnight in SM broth (DSMZ medium 7) at 28°C. DNA was isolated using the Genomic 500 DNA kit (Qiagen, Hilden, Germany) following the instructions of the manufacturer. Based on its phylogenetic relationship to *Spirosoma linguale* (2), the cell lysis modification procedure and prolonged incubation time for that bacterium were applied as well.

The genome was sequenced at the Functional Genomics Center Zurich (Zurich, Switzerland) with Roche 454 GS FLX Titanium chemistry. Two libraries were prepared: half a plate with an 8-kb insert paired-end library and half a plate with a shotgun library. This strategy yielded 1,325,215 reads with an average size of 378 nucleotides (nt) for the shotgun library and 330 nt for the paired-end library. Both sets of reads were assembled using the GS DeNovo Assembler version 2.5.3 (Roche) with various parameter settings. The two best assemblies were merged using Minimus 2 version 3.00 (5) and scaffolded using Oslay software version 1.0 (4). The assembly was further improved by repeatedly mapping the paired-end reads against the assembly and manually checking the gap regions, which allowed the closure of most gaps. The final genome sequence of *F. limi* BUZ 3^T is composed of 11 scaffolded contigs representing the chromosome (N₅₀, 2,007,105), a large plasmid (pFLIM01), and two small plasmids (pFLIM02 and pFLIM03). The total size of the chromosomal contigs is 7,454,326 bp, with a G+C content of 52.7%. The size of pFLIM01 is 151,932 bp with a G+C content of 49.8%, while the two small plasmids are 14,391 bp (pFLIM02; G+C content, 47.03%) and 8,321 bp (pFLIM03; G+C content, 37.4%). The average coverage ranged between 38.3-fold (pFLIM03) and 58.6-fold (pFLIM01), with the chromosome having 44.4-fold average coverage.

The GenDB annotation platform (3) was employed for the genome annotation. A total of 6,710 protein-coding genes (CDSs) and 49 structural RNAs on the chromosome and 197 CDSs on the three plasmids were annotated.

Members of the phylum *Bacteroidetes* are widely distributed in aquatic environments and have several interesting metabolic (e.g., degradation of complex polymers) and morphological (e.g., multicellular filamentation) features. Therefore, increasing the number of genomes sequenced for this phylum will improve knowledge of these organisms. A comprehensive comparative genome analysis is under way.

Nucleotide sequence accession numbers. The 11 contigs of the chromosome of *F. limi* BUZ 3^T were deposited at EBI under the accession numbers CAIT01000001 through CAIT01000011 (project number CAIT01000000), and the plasmids were deposited under accession numbers HE805916 (pFLIM01), HE805917 (pFLIM02), and HE805918 (pFLIM03).

ACKNOWLEDGMENTS

This project was funded by the University of Zurich and the Kanton of Zurich, Switzerland.

We are thankful to C. Aquino and R. Lecca for library preparation. We thank the Institute for Genome Sciences (IGS; University of Maryland), which performed the annotation of the first draft of the genome sequence.

REFERENCES

1. Filippini M, Kaech A, Ziegler U, Bagheri HC. 2011. *Fibrisoma limi* gen. nov., sp. nov., a filamentous bacterium isolated from tidal flats. *Int. J. Syst. Evol. Microbiol.* 61:1418–1424.
2. Lail K, et al. 2010. Complete genome sequence of *Spirosoma linguale* type strain (1T). *Stand. Genomic Sci.* 2:176–185.
3. Meyer F, et al. 2003. GenDB: an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res.* 31:2187–2195.
4. Richter DC, Schuster SC, Huson DH. 2007. OSLay: optimal syntenic layout of unfinished assemblies. *Bioinformatics* 23:1573–1579.
5. Schatz MC, et al. 23 December 2011, posting date. Hawkeye and AMOS: visualizing and assessing the quality of genome assemblies. *Brief. Bioinform.* doi:10.1093/bib/bbr074.

Received 17 May 2012 Accepted 31 May 2012

Address correspondence to Manuela Filippini, manuela.filippini@ieu.uzh.ch.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00869-12